

THE GROWTH OF ENTIRE FUNCTIONS OF GENUS ZERO

DANG DUC TRONG AND TRUONG TRUNG TUYEN

ABSTRACT. In this paper we shall consider the asymptotic growth of $|P_n(z)|^{1/k_n}$ where $P_n(z)$ is a sequence of entire functions of genus zero. Our results extend a result of J. Muller and A. Yavrian. We shall prove that if the sequence of entire functions has a geometric growth at each point in a set E being non-thin at ∞ then it has a geometric growth in $\mathbb{C}$ also. Moreover, if E has some more properties, a similar result also holds for a more general kind of growth. Even in the case where P_n are polynomials, our results are new in the sense that it does not require $k_n \succeq \deg(P_n)$ as usually required.

1. INTRODUCTION AND MAIN RESULTS.

The growth at infinity of entire functions is a topic of great concernment. In [3], the authors gained interested results which combine the growth of a sequence of polynomials on a "small subset" of $\mathbb{C}$ with the growth of itself on the whole plane. The "small subsets" as mentioned are non-thin. We recall that (see [3]) a domain G , with ∂G having positive capacity, is non-thin at an its boundary point $\zeta \in \partial G$ (or ζ is a regular point of G) if and only if

$$\lim_{z \in G, z \rightarrow \zeta} g(z, w) = 0$$

for all $w \in G$ where $g(.,.)$ is the Green function of G . One of the main results in [3] is stated as below (see Lemma 2 in [3])

Proposition 1. *Let (d_n) be a sequence of positive numbers and let (P_n) be a sequence of polynomials satisfying $\deg(P_n) \leq d_n$. If $E \subseteq \mathbb{C}$ is closed and non-thin at ∞ so that*

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/d_n} \leq 1, \quad \text{for all } z \in E,$$

then

$$\limsup_{n \rightarrow \infty} \|P_n\|_R^{1/d_n} \leq 1, \quad \text{for all } R > 0,$$

where $\|P_n\|_R = \sup\{|P_n(z)| : |z| \leq R\}$.

Date: February 2, 2008.

2000 Mathematics Subject Classification. 30C85, 30D15, 31A15.

Key words and phrases. Capacity; Entire functions of genus zero; Geometric growth; Non-thin set.

Saying roughly, Proposition 1 states that if E is a set being non-thin at infinity and (P_n) is a sequence of polynomials having a geometric growth at each point in E then (P_n) has a geometric growth in $\mathcal{C}$ also.

There are two interesting questions rising from this result

- 1) Does the conclusion of Proposition 1 still hold if P_n are non-polynomial entire functions?
- 2) If (P_n) has a non-geometric growth in E , i.e., if instead of the condition

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/d_n} \leq 1, \text{ for all } z \in E,$$

we requires only that

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/d_n} \leq h(|z|), \text{ for all } z \in E,$$

where h is not necessary a bounded function, does (P_n) remain the same rate of growth in $\mathcal{C}$?

To answer both these questions, a class of entire functions seeming appropriate is the class of entire functions of genus zero. On a hand, this class is similar to the class of polynomials. A function P of this class can be factorized by monomials whose roots are roots of P . On the other hand, this class is fairly wide. It is very close to entire functions of class Cartwright (see [4] for definition and properties of this class) containing entire functions generated from Fourier's transformations.

For answering Question 1, for an entire function P of genus zero, we define a degree $d^*(P)$ similar to the degree of polynomials, which satisfies $d^*(P) \leq d(P)$ if P is a polynomial where $d(P)$ is the ordinary degree of P . In fact, we obtain a nearly complete answer for the question: what is the necessary and sufficient condition under which a sequence of entire functions of genus zero has a geometric growth?

For Question 2, the answer is confirmation in the case $h(z)$ is a polynomial and E is a closed set satisfying

$$\limsup_{R \rightarrow \infty} \frac{\log \text{cap}(E_R)}{\log R} = \beta > 0,$$

where

$$E_R = E \cap \{z : |z| \leq R\}.$$

In proving this result we don't use the property that E is non-thin at infinity. So from this result we immediately get that E must be non-thin at infinity. As known in [3], the authors showed that such E 's sets are non-thin at infinity by using Wiener's criterion.

This paper consists of four parts. In Section 2 we shall set some notations and state (and prove) some necessary lemmas. In Section 3 we prove two results in which Theorem 1 can be seen as a direct generalization of Proposition 1. The results in this Section consist of a sufficient and nearly-necessary condition under which a sequence of entire functions of genus zero will have

a geometric growth, so answer Question 1. In Section 4 we prove a result answering Question 2.

2. NOTATIONS AND LEMMAS

For an entire function f we use notations

$$\begin{aligned} \|f\|_R &= \sup_{\{|z| \leq R\}} |f(z)| \\ C(f, R) &= \exp\left\{\frac{1}{2\pi} \int_0^{2\pi} \log |f(Re^{it})| dt\right\}, \\ \eta(f, R) &= \text{the number of elements of } \{z : 0 < |z| \leq R, f(z) = 0\}. \end{aligned}$$

An entire function is called of genus zero if its order is less than 1. We recall that (see Lecture 1 in [4]): if

$$P(z) = \sum_{n=0}^{\infty} a_n z^n$$

then its order ρ can be written as

$$\rho = \limsup_{n \rightarrow \infty} \frac{n \log n}{\log(1/|a_n|)}.$$

If P is of genus zero then P can be expressed as follows (see §4.2 in [4])

$$P(z) = az^\alpha \prod_j (1 - z/z_j),$$

where $\alpha \in \mathbf{N}$ and z_j 's are non-zero complex numbers satisfying

$$\sum_j 1/|z_j| < \infty.$$

If P is of genus zero and P is expressed as above, we define

$$d^*(P) = \alpha + \sum_{|z_j| \leq 1} 1 + \sum_{|z_j| > 1} 1/|z_j|,$$

and we shall call it the "degree" of the entire function P . In case of P polynomials, one has $d^*(P) \leq d(P)$.

Hereafter we always consider a sequence of positive numbers (k_n) and a sequence of entire functions of genus zero (P_n) having the following form

$$P_n(z) = a_n z^{\alpha_n} \prod_j (1 - z/z_{n,j}),$$

and such that $k_n \geq d^*(P_n)$ for all $n \in \mathbf{N}$.

We put

$$\begin{aligned} C_0 &= \limsup_{n \rightarrow \infty} C(P_n, 1)^{1/k_n}, \\ C_0^* &= \liminf_{n \rightarrow \infty} C(P_n, 1)^{1/k_n}, \\ \eta(R) &= \limsup_{n \rightarrow \infty} \frac{\eta(P_n, R)}{k_n}. \end{aligned}$$

For definition of capacity of a compact set, its properties and its relations to the Green's function and the harmonic measure of the set, one can refer to [2].

Lemma 1. (i) If $C_0 = 0$ then for all $R > 0$

$$\limsup_{n \rightarrow \infty} \|P_n\|_R^{1/k_n} = 0.$$

(ii) Assume that $C_0^* > 0$ and

$$\limsup_{n \rightarrow \infty} C(P_n, R)^{1/k_n} \leq h(R),$$

where h satisfies

$$\liminf_{R \rightarrow \infty} \frac{\log h(R)}{\log R} \leq \tau.$$

Then for all $R > 0$ we have $\eta(R) \leq \tau$.

Proof. (i) For each $n \in \mathbb{N}$, $z \in \mathbb{C}$, by applying Jensen's identity (see, e.g., Theorem 15.18 in [5]) gives

$$\begin{aligned} |P_n(z)| &\leq |a_n| |z|^{\alpha_n} \prod_j (1 + R/|z_{n,j}|) \\ &\leq |a_n| |z|^{\alpha_n} \prod_{|z_{n,j}| \leq 1} (1 + R)/|z_{n,j}| \exp\{|z| \sum_{|z_{n,j}| > 1} 1/|z_{n,j}|\} \\ &= (|a_n| |z|^{\alpha_n} \prod_{|z_{n,j}| \leq 1} 1/|z_{n,j}|) (1 + R)^{\eta(P_n, 1)} \exp\{|z| \sum_{|z_{n,j}| > 1} 1/|z_{n,j}|\} \\ &= C(P_n, 1) (1 + R)^{\eta(P_n, 1)} \exp\{|z| \sum_{|z_{n,j}| > 1} 1/|z_{n,j}|\} \\ &\leq C(P_n, 1) (1 + R)^{d^*(P_n)} \exp\{|z| d^*(P_n)\}. \end{aligned}$$

Thus

$$\begin{aligned} \limsup_{n \rightarrow \infty} \|P_n\|_R^{1/k_n} &\leq \limsup_{n \rightarrow \infty} C(P_n, 1)^{1/k_n} (1 + R)^{d^*(P_n)/k_n} \exp\{|z| d^*(P_n)/k_n\} \\ &\leq (1 + R) \exp\{R\} \limsup_{n \rightarrow \infty} C(P_n, 1)^{1/k_n} \\ &= (1 + R) \exp\{R\} C_0 = 0. \end{aligned}$$

(ii) Fix $R > 0$ and choose $s > R$. Applying Jensen's formula gives

$$C(P_n, R) = |a_n| R^{\alpha_n} \prod_j \max\{1, \frac{R}{|z_{n,j}|}\}.$$

We get

$$C(P_n, s) \geq C(P_n, R)(s/R)^{\eta(P_n, R)},$$

hence

$$\begin{aligned} h(s) &\geq \limsup_{n \rightarrow \infty} C(P_n, s)^{1/k_n} \\ &\geq \limsup_{n \rightarrow \infty} C(P_n, R)^{1/k_n} (s/R)^{\eta(P_n, R)/k_n} \\ &\geq \liminf_{n \rightarrow \infty} C(P_n, R)^{1/k_n} \limsup_{n \rightarrow \infty} (s/R)^{\eta(P_n, R)/k_n} \\ &= C_0^*(s/R)^{\eta(R)}. \end{aligned}$$

Thus

$$\begin{aligned} \tau &\geq \liminf_{s \rightarrow \infty} \frac{\log h(R)}{\log R} \\ &\geq \liminf_{s \rightarrow \infty} \frac{\log C_0^*(s/R)^{\eta(R)}}{\log s} \\ &= \eta(R). \end{aligned}$$

This completes the proof of Lemma 1. $\square$

We end this section with a result relating the maximum and logarithm norms

Lemma 2. *Assume that $C_0 < \infty$, and that*

$$(2.1) \quad \lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{\left| \sum_{|z_{n,j}| \geq R} \frac{1}{z_{n,j}} \right|}{k_n} = 0,$$

and there exists a sequence (R_n) of positive real numbers tending to ∞ such that

$$(2.2) \quad \limsup_{n \rightarrow \infty} \frac{\eta(P_n, R_n)}{k_n} < \infty.$$

If

$$\liminf_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{\log C(P_n, R)^{1/k_n}}{\log R} \leq \tau,$$

then for all $R > 0$ we have

$$\limsup_{n \rightarrow \infty} \|P_n\|_R^{1/k_n} \leq C_0(1+R)^\tau.$$

Proof. Without loss of generality, we may assume that $R > 1$.

In view of Lemma 1 (i) we only consider the case in which $C_0 > 0$.

To prove the result we need to show that: each subsequence of (P_n) contains a subsequence that satisfies the conclusion of Lemma 2. We choose a subsequence, still denoted by (P_n) , such that

$$\lim_{n \rightarrow \infty} C(P_n, 1)^{1/k_n} = C_1.$$

We note that $C_1 \leq C_0$.

If $C_1 = 0$ then we get the conclusion by Lemma 1 (i).

If $C_1 > 0$ then applying Lemma 1 for this subsequence gives $\eta(R) \leq \tau$, for all $R > 0$. Choosing a $\beta > 1$, we have

$$\begin{aligned} \|P_n\|_R &\leq |a_n| R^{\alpha_n} \prod_{|z_{n,j}| \leq \beta R} (1 + R/|z_{n,j}|) \sup_{|z| \leq R} \prod_{|z_{n,j}| > \beta R} |1 - z/z_{n,j}| \\ &\leq C(P_n, 1)(1 + R)^{\eta(P_n, 1)} \times \prod_{1 < |z_{n,j}| \leq \beta R} (1 + R/|z_{n,j}|) \times \\ &\quad \times \sup_{|z| \leq R} \exp\left\{ \sum_{|z_{n,j}| \geq \beta R} |1 - z/z_{n,j}| - 1 \right\} \\ &\leq C(P_n, 1)(1 + R)^{\eta(P_n, \beta R)} \sup_{|z| \leq R} \exp\left\{ \sum_{|z_{n,j}| \geq \beta R} |1 - z/z_{n,j}| - 1 \right\}. \end{aligned}$$

Noting that for $|R/z_{n,j}| \leq 1/\beta$, we have

$$\begin{aligned} |1 - z/z_{n,j}| - 1 &= \frac{|1 - z/z_{n,j}|^2 - 1}{|1 - z/z_{n,j}| + 1} \\ &= \frac{-z/z_{n,j} - \bar{z}/\bar{z}_{n,j} + R^2/|z_{n,j}|^2}{|1 - z/z_{n,j}| + 1} \\ &\leq \max\left\{ \frac{-z/z_{n,j} - \bar{z}/\bar{z}_{n,j} + R/(\beta|z_{n,j}|)}{2 - 1/\beta}, \right. \\ &\quad \left. \frac{-z/z_{n,j} - \bar{z}/\bar{z}_{n,j} + R/(\beta|z_{n,j}|)}{2} \right\} \\ &\sim \frac{-z/z_{n,j} - \bar{z}/\bar{z}_{n,j} + R/(\beta|z_{n,j}|)}{2}, \end{aligned}$$

with β large enough. Hence

$$\begin{aligned} \|P_n\|_R &\leq C(P_n, 1)(1 + R)^{\eta(P_n, \beta R)} \times \\ &\quad \times \sup_{|z| \leq R} \exp\left\{ \frac{-\sum_{|z_{n,j}| \geq \beta R} (z/z_{n,j} + \bar{z}/\bar{z}_{n,j}) + (R/\beta) \sum_{|z_{n,j}| \geq \beta R} 1/|z_{n,j}|}{2} \right\} \\ &\leq C(P_n, 1)(1 + R)^{\eta(P_n, \beta R)} \times \\ &\quad \times \exp\left\{ \frac{2R|\sum_{|z_{n,j}| \geq \beta R} 1/|z_{n,j}| + (R/\beta) \sum_{|z_{n,j}| \geq \beta R} 1/|z_{n,j}|}{2} \right\}. \end{aligned}$$

It follows that

$$\begin{aligned}
\limsup_{n \rightarrow \infty} \|P_n\|_R^{1/k_n} &\leq C_1(1+R)^{\eta(R)} \times \\
&\quad \times \limsup_{n \rightarrow \infty} \exp\left\{R \frac{|\sum_{|z_{n,j}| \geq \beta R} 1/z_{n,j}| + (1/\beta) \sum_{|z_{n,j}| \geq \beta R} 1/|z_{n,j}|}{2k_n}\right\} \\
&\leq C_0(1+R)^\tau \times \\
&\quad \times \limsup_{n \rightarrow \infty} \exp\left\{2R \frac{|\sum_{|z_{n,j}| \geq \beta R} 1/z_{n,j}| + (1/\beta) \sum_{|z_{n,j}| \geq \beta R} 1/|z_{n,j}|}{k_n}\right\}.
\end{aligned}$$

Letting β tend to ∞ we get

$$\begin{aligned}
\limsup_{n \rightarrow \infty} \|P_n\|_R^{1/k_n} &\leq C_0(1+R)^\tau \lim_{\beta \rightarrow \infty} \exp\left\{R \frac{|\sum_{|z_{n,j}| \geq \beta R} 1/z_{n,j}|}{k_n}\right\} \\
&= C_0(1+R)^\tau.
\end{aligned}$$

□

3. THE CASE OF GEOMETRIC GROWTH

Theorem 1. *Let E be a closed set being non-thin at ∞ . Assume that*

$$(3.1) \quad \lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{|\sum_{|z_{n,j}| \geq R} 1/z_{n,j}|}{k_n} = 0,$$

and there exists a sequence $\{R_n\}$ of positive real numbers tending to ∞ such that

$$(3.2) \quad \limsup_{n \rightarrow \infty} \frac{\eta(P_n, R_n)}{k_n} < \infty.$$

If for each $z \in E$ one has

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1,$$

then

$$\limsup_{n \rightarrow \infty} \|P_n\|_R^{1/k_n} \leq 1, \text{ for all } R > 0.$$

As will be shown in Theorem 2, from the assumptions of Lemma 1 (ii), we shall obtain conditions (3.1) and (3.2)

Theorem 1 generalizes Proposition 1. To show this end, we note that if P_n are polynomials and $d_n \geq \deg(P_n)$ then

$$\lim_{R \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{|\sum_{|z_{n,j}| \geq R} 1/z_{n,j}|}{d_n} \leq \lim_{R \rightarrow \infty} \frac{1}{R} = 0,$$

and for all $n \in \mathbb{N}$ and $R > 0$

$$\frac{\eta(P_n, R)}{d_n} \leq 1.$$

Proof. For each compact set A in the complex plane we take $g(A, z)$ to be its Green's function having pole at infinity of the unbounded component of $\mathbb{C} \setminus A$ and extend $g(A, z)$ to be zero outside that component.

For each $R > 0$ let E_R^* be the union of E_R with the bounded components of $\mathbb{C} \setminus E_R$. Then $E_R \subset E_R^*$ and $\mathbb{C} \setminus E_R^*$ has no bounded components. For $z \in E_R^*$ we claim that

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1.$$

In fact, the compare principle gives $g(E_R^*, z) \leq g(E_R, z)$ for all $z \in \mathbb{C}$. Thus by Lemma 2 in [3], for all $s > 0$ we have

$$\lim_{R \rightarrow \infty} \int_0^{2\pi} g(E_R^*, se^{it}) dt \leq \lim_{R \rightarrow \infty} \int_0^{2\pi} g(E_R, se^{it}) dt = 0.$$

Arguing as in Step 1 in proof of Lemma 2 in [3], we have

$$(3.3) \quad \limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq 1, \text{ for each } z \in E_R^*.$$

For each $n \in \mathbb{N}$ we put

$$Q_n(z) = a_n z^{\alpha_n} \prod_{|z_{n,j}| \leq R_n} (1 - z/z_{n,j}).$$

By (3.1) we have

$$\limsup_{n \rightarrow \infty} |Q_n(z)|^{1/k_n} = \limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n},$$

for all $z \in \mathbb{C}$. Indeed, putting $P_n(z) = Q_n(z)H_n(z)$, to prove this assertion we need only to prove that

$$\lim_{n \rightarrow \infty} |H_n(z)|^{1/k_n} = 1,$$

for any fixed $z \in \mathbb{C}$ where

$$H_n(z) = \prod_{|z_{n,j}| > R_n} (1 - z/z_{n,j}).$$

Arguing as in the proof of Lemma 2, using (3.1) we get that

$$\limsup_{n \rightarrow \infty} |H_n(z)|^{1/k_n} \leq 1.$$

By the Taylor's expansion for the function $\log(1 + \epsilon)$ for $|\epsilon|$ small enough, we see that $t \geq \exp\{t - 1 - 2(1 - t)^2\}$ for t near 1. Since z is fixed, for n large enough we have $|1 - z/z_{n,j}|$ is near 1 if $|z_{n,j}| \geq R_n$, so using the same argument in the proof of Lemma 2 we have

$$\begin{aligned} |1 - z/z_{n,j}| &\geq \exp\{|1 - z/z_{n,j}| - 1 - 2(|1 - z/z_{n,j}|)^2\} \\ &\geq \exp\left\{\frac{-z/z_{n,j} - \bar{z}/\bar{z}_{n,j} - 3|z|^2/|z_{n,j}|^2}{|1 - z/z_{n,j}| + 1}\right\}. \end{aligned}$$

Thus similarly we have

$$\liminf_{n \rightarrow \infty} |H_n(z)|^{1/k_n} \geq 1.$$

Combining above results we get that

$$\lim_{n \rightarrow \infty} |H_n(z)|^{1/k_n} = 1.$$

By Lebesgue's dominated convergence theorem we get

$$\limsup_{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \log |Q_n(se^{it})|^{1/k_n} dt = \limsup_{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \log |P_n(se^{it})|^{1/k_n} dt,$$

for all $s > 0$.

From (3.3) we have

$$\limsup_{n \rightarrow \infty} |Q_n(z)|^{1/k_n} \leq 1, \text{ for all } z \in E_R^*.$$

Applying Bernstein's inequality (see e.g., [3]) to polynomials $Q'_n s$ and arguing as in Step 2 in the proof of Lemma 2 in [3] we get

$$\limsup_{n \rightarrow \infty} \log |Q_n(z)|^{1/k_n} \leq \kappa g(E_R^*, z), \text{ for all } z \in C,$$

where

$$\kappa = \limsup_{n \rightarrow \infty} \frac{\eta(P_n, R_n)}{k_n} < \infty.$$

Fixing $s > 0$, intergrating above inequality on the circle $|z| = s$, applying Fatou's Lemma (see e.g., Lemma 1.28 in [5]), and letting $R \rightarrow \infty$ we get

$$\limsup_{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \log |Q_n(se^{it})|^{1/k_n} dt \leq 0.$$

Thus

$$\limsup_{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \log |P_n(se^{it})|^{1/k_n} dt \leq 0,$$

for all $s > 0$, which gives

$$\limsup_{n \rightarrow \infty} \|P_n\|_R^{1/k_n} \leq 1,$$

for all $R > 0$, by Lemma 2. $\square$

Theorem 2. Assume that $C_0^* > 0$. Given a function $h : (0, +\infty) \rightarrow \mathbb{R}$ satisfying

$$\limsup_{R \rightarrow \infty} \frac{\log h(R)}{\log R} < \infty.$$

If for all $R > 0$

$$\limsup_{n \rightarrow \infty} C(P_n, R)^{1/k_n} \leq h(R),$$

then there exists a sequence of positive numbers $R_n \rightarrow \infty$ such that

$$\limsup_{n \rightarrow \infty} \frac{|\sum_{|z_{n,j}| > R_n} 1/z_{n,j}|}{k_n} = 0$$

$$\limsup_{n \rightarrow \infty} \frac{\eta(P_n, R_n)}{k_n} < \infty.$$

We recall that the conclusions of Theorem 2 are equivalent to the conditions (3.1) and (3.2).

Proof. By Lemma 1 we get

$$\limsup_{n \rightarrow \infty} \eta(R) \leq \tau, \text{ for all } R > 0.$$

For each $s \in \mathbf{N}$ we choose $n_s \geq s$ such that $\eta(P_n, s)/k_n \leq \tau + 1/s$ for all $n \geq n_s$. Moreover it can be taken such that $n_1 < n_2 < \dots$. We put $R_n = s$ if $n_s \leq n < n_{s+1}$. We prove that these R_n are the desired sequence.

For each $n \in \mathbf{N}$ we put

$$Q_n(z) = a_n z^{\alpha_n} \prod_{|z_{n,j}| \leq R_n} (1 - z/z_{n,j}),$$

and

$$H_n(z) = \prod_{|z_{n,j}| \geq R_n} (1 - z/z_{n,j}).$$

We have

$$\liminf_{n \rightarrow \infty} C(Q_n, 1)^{1/k_n} = \liminf_{n \rightarrow \infty} C(P_n, 1)^{1/k_n} = C_0^*,$$

and that the sequence $\{Q_n, k_n\}$ satisfies the conditions (3.1) and (3.2) of Theorem 1. It follows that for all $t \in [0, 2\pi]$ and $R > 0$, there exists $s > R$ such that

$$\limsup_{n \rightarrow \infty} |Q_n(se^{it})|^{1/k_n} \geq C_0^*/2,$$

since the sets $E_t = \{z : z = se^{it}, s \geq R\}$ is closed and non-thin at ∞ .

To prove the other conclusion of Theorem we need to show that each subsequence of (P_n) has a subsequence, still denoted by (P_n) , to which the conclusion is satisfied. We have

$$k_n \geq d^*(P_n) \geq \sum_{|z_{n,j}| \geq 1} 1/|z_{n,j}|, \text{ for all } n \in \mathbf{N},$$

so we can assume that

$$\lim_{n \rightarrow \infty} \frac{\sum_{|z_{n,j}| > R_n} 1/z_{n,j}}{k_n} = \alpha.$$

Let $-\pi < \theta \leq \pi$ be such that $\alpha = |\alpha|e^{i\theta}$.

For each $R > 0$ we choose $s > R$ such that

$$\limsup_{n \rightarrow \infty} |Q_n(-se^{-i\theta})|^{1/k_n} \geq C_0^*/2.$$

Noting that $|1 - z/z_{n,j}| \geq \exp\{|1 - z/z_{n,j}| - 1 - 2(|1 - z/z_{n,j}| - 1)^2\}$, and arguing as in the proof of Lemma 1 (ii) and Theorem 1 we get

$$\liminf_{n \rightarrow \infty} |H_n(-se^{-i\theta})|^{1/k_n} \geq \exp\{\limsup_{n \rightarrow \infty} s \frac{|\sum_{|z_{n,j}| > R_n} 1/z_{n,j}|}{k_n}\},$$

thus

$$\begin{aligned} h(s) &\geq \limsup_{n \rightarrow \infty} ||P_n||_s^{1/k_n} \\ &\geq \limsup_{n \rightarrow \infty} |P_n(-se^{i\theta})|_s^{1/k_n} \\ &\geq \limsup_{n \rightarrow \infty} |Q_n(-se^{-i\theta})|^{1/k_n} \liminf_{n \rightarrow \infty} |H_n(-se^{-i\theta})|^{1/k_n} \\ &\geq C_0^*/2 \exp\{\limsup_{n \rightarrow \infty} s \frac{|\sum_{|z_{n,j}| > R_n} 1/z_{n,j}|}{k_n}\}. \end{aligned}$$

From the properties of h we get the conclusion of Theorem 2. $\square$

4. THE CASE OF NON-GEOMETRIC GROWTH

Theorem 3. *Let E be a closed set such that*

$$(4.1) \quad \limsup_{R \rightarrow \infty} \frac{\log \text{cap}(E_R)}{\log R} = \beta > 0.$$

Assume that (3.1) and (3.2) hold. If for all $z \in E$ we have

$$\limsup_{n \rightarrow \infty} |P_n(z)|^{1/k_n} \leq h(|z|),$$

where

$$\limsup_{R \rightarrow \infty} \frac{\log h(R)}{\log R} \leq \gamma < \infty.$$

Then for all $R > 0$ we have

$$\limsup_{n \rightarrow \infty} ||P_n||_R^{1/k_n} \leq C_0(1 + R)^{\gamma/\beta}.$$

Proof. Take $s_n \rightarrow \infty$ such that

$$\lim_{n \rightarrow \infty} \frac{\log \text{cap}(E_{s_n})}{\log s_n} = \beta.$$

We can assume that $E \cap \{z : |z| = s_n\} = \emptyset$ for all $n \in \mathbb{N}$. Indeed, we can choose s'_n such that

$$\lim_{n \rightarrow \infty} \frac{\log s_n}{\log s_{n+1}} = \lim_{n \rightarrow \infty} \frac{\log n}{\log s_n} = 0.$$

Let $E^* = E \setminus \bigcup_{n=1}^{\infty} \{z : s_n < |z| < 1 + s_n\}$. Then E^* is closed, and for all $n \in \mathbb{N}$ we have

$$\text{cap}(E_{s_n}^*) \geq \text{cap}(E_{s_n}) - \text{cap}(E_{s_{n-1}}) \geq \text{cap}(E_{s_n}) - \log s_{n-1},$$

hence

$$\lim_{n \rightarrow \infty} \frac{\log \operatorname{cap}(E_{s_n}^*)}{\log s_n} = \beta.$$

Replacing E and s_n by E^* and $s_n + \frac{1}{2}$, we see that the conditions of Theorem still hold and $E \cap \{z : |z| = s_n\} = \emptyset$ for all $n \in \mathbf{N}$. It follows that (see e.g., formula (8.3) page 114 in [2])

$$g(E_{s_n}, s_n e^{it}) = -\log \operatorname{cap}(E_{s_n}) + \int_{\partial E_{s_n}} \log |s_n e^{it} - \zeta| d\mu(E_{s_n}, \zeta),$$

where $n \in \mathbf{N}$, $t \in [0, 2\pi]$ and $\mu(., .)$ is the harmonic measure. Integrating this identity and applying Fubini's Theorem (see e.g., Theorem 7.8 in [5]) we get

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} g(E_{s_n}, s_n e^{it}) dt &= -\log \operatorname{cap}(E_{s_n}) \\ &\quad + \frac{1}{2\pi} \int_0^{2\pi} \int_{\partial E_{s_n}} \log |s_n e^{it} - \zeta| d\mu(E_{s_n}, \zeta) \\ &= -\log \operatorname{cap}(E_{s_n}) \\ &\quad + \int_{\partial E_{s_n}} d\mu(E_{s_n}, \zeta) \frac{1}{2\pi} \int_0^{2\pi} \log |s_n e^{it} - \zeta| dt \\ &= -\log \operatorname{cap}(E_{s_n}) + \int_{\partial E_{s_n}} d\mu(E_{s_n}, \zeta) \log s_n \\ &= \log s_n - \log \operatorname{cap}(E_{s_n}), \end{aligned}$$

where we have used

$$\begin{aligned} \frac{1}{2\pi} \int_0^{2\pi} \log |s_n e^{it} - \zeta| dt &= \max\{\log s_n, \zeta\} = \log s_n, \\ \int_{\partial E_{s_n}} d\mu(E_{s_n}, \zeta) &= 1. \end{aligned}$$

Hence we have

$$\lim_{n \rightarrow \infty} \frac{1}{2\pi} \int_0^{2\pi} \frac{g(E_{s_n}^*, s_n e^{it})}{\log s_n} dt = 1 - \beta.$$

Put, as in Theorem 1,

$$\kappa = \limsup_{n \rightarrow \infty} \frac{\eta(P_n, R_n)}{k_n}.$$

Arguing as in the proof of Theorem 1 we get

$$\limsup_{n \rightarrow \infty} \log |Q_n(z)|^{1/k_n} \leq \log h(R) + \kappa g(E_R, z),$$

for all $z \in \mathcal{C}$ and $R > 0$ such that $\operatorname{cap}(E_R) > 0$.

It follows that in view of the definition of $C(P, R)$ that

$$\limsup_{n \rightarrow \infty} \log C(P_n, s_m)^{1/k_n} / \log s_m \leq \log h(s_m) / \log s_m + \kappa \frac{1}{2\pi} \int_0^{2\pi} \frac{g(E_{s_m}^*, s_m e^{it})}{\log s_m},$$

for all $m \in N$. Hence

$$\begin{aligned} \liminf_{s \rightarrow \infty} \limsup_{n \rightarrow \infty} \log C_n(P_n, s)^{1/k_n} / \log s &\leq \limsup_{m \rightarrow \infty} [\log h(s_m) / \log s_m \\ &\quad + \kappa \frac{1}{2\pi} \int_0^{2\pi} \frac{g(E_{s_m}^*, s_m e^{it})}{\log s_m}] \\ &\leq \gamma + \kappa(1 - \beta). \end{aligned}$$

By Lemma 1 we have

$$\liminf_{s \rightarrow \infty} \limsup_{n \rightarrow \infty} \log C_n(P_n, s)^{1/k_n} / \log s \geq \kappa.$$

Thus

$$\kappa \leq \gamma + \kappa(1 - \beta),$$

or

$$\kappa \leq \gamma / \beta.$$

Hence

$$\liminf_{s \rightarrow \infty} \limsup_{n \rightarrow \infty} \frac{\log C(P_n, s)^{1/k_n}}{\log s} \leq \gamma + \frac{\gamma}{\beta}(1 - \beta) = \frac{\gamma}{\beta}.$$

Applying Lemma 2 we get the conclusion of Theorem 3. $\square$

REFERENCES

- [1] Peter Borwein and Tamas Erdeleyi, *Polynomials and polynomial inequalities*, Springer - Verlag, New York, 1995.
- [2] W. H. J. Fuchs, *Topics in the theory of functions of one complex variable*, D. Van Nostrand Company Inc., Princeton-New Jersey-Toronto-London-Melbourne, 1967.
- [3] J. Muller and A. Yavrian, *On polynomials sequences with restricted growth near infinity*, Bull. London Math. Soc. **34** 2002, 189–199.
- [4] B. Ya. Levin, *Lectures on entire functions*, Translations of Mathematical Monographs **150**, AMS, Providence, Rhode Island, 1996.
- [5] Walter Rudin, *Real and complex analysis*, Mladinska Knjiga, Ljubljana, 1970.

DEPARTMENT OF MATHEMATICS, HOCHIMINH CITY NATIONAL UNIVERSITY, 227 NGUYEN VAN CU, Q5, HOCHIMINH CITY, VIETNAM

E-mail address: ddtrong@mathdep.hcmuns.edu.vn

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, RAWLES HALL, BLOOMINGTON, IN 47405

E-mail address: truongt@indiana.edu